PTO/SB/08b (06-03) Approved for use through 06/30/2003. OMB 0651-0031
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

eduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Substitute for form 1	449B/PTO					Complete if Known	
				Ī	Application Number	10/076,708	
INFORM	ATION DI	SCLC	SURE		Filing Date	February 15, 2002	
STATEM	ENT BY	APPLI	CANT	Ì	First Named Inventor	Satish K. Sharma	
					Art Unit	1646	
(Use as many sheets as necessary)					Examiner Name	Chernyshev, Olga N	
Sheet	1	of		1	Attorney Docket Number	6322.N	

		NON PATENT LITERATURE DOCUMENTS	
Examiner Initials*	Cite No.1	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	T²
	1	GOETZ, J ET AL: "Formation of neurofibrillary tangles in P301L tau transgenic mice induced by Abeta42 fibrilis" Science, American Association for the Advancement of Science, US, vol. 293, no. 5534, pages 1491-1495, 2001.	
	2	GOURAS GUNNAR K ET AL: "Intraneuronal Abeta 42 accumulation in human brain" American Journal of Pathology, vol. 156, no. 1, pages 15-20, January 2000.	
***************************************	3	HARDY JOHN ET AL: "Genetic dissection of Alzheimer's disease and related demetias: Amyloid and its relationship to tau" Nature Neuroscience, vol. 1, no. 5, pages 355-358, 1998.	
	4	LEWIS J ET AL: "Enhanced neurofibrillary degeneration in transgenic mice expressin mutant tau and APP" Science, American Association for the Advancement of Science, vol. 293, no. 5534, pages 1487-1491, 2001.	
***************************************	5	RANK KB ET AL: "Direct interaction of soluble human recombinant tau protein with Abeta 1-42 results in tau aggregation and hperphosphorylation by tau protein kinase II" Febs Letters, Elsevier Science Publishers, Amsterdam, NL, vol. 514, no. 2-3, pages 263-268, 2002.	
	6	TAKASHIMA A ET AL: "Amyloid beta peptide (25-35) induces tau phosphorylation and decrease microtubule-forming ability in rat hippocampal culture" Abstracts of the Society for Neuroscience, WA, DC US, vo. 21, no. 1-3, page 1719, 1995.	
	7	TOMIDOKORO YASUSHI ET AL: "Abeta amyloidosis induces the initial stage of tau accumulation in APPSW mice" Neuroscience Letters, vol. 299, no. 3, pages 169-172, 2001.	
***************************************	***************************************		!
·			

Examiner	Date	
Signature	 Considered	

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Applicant's unique citation designation number (optional). 2 Applicant is to place a check mark here if English language Translation is attached.

This collection of information is required by 37 CFR 1.97 and 1.98. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 2 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

f. Info disi

LIST	FRE	ERENCES CI	TED BY APPLIC	ANT	Atty	Docket No. 6322.N			Serial N 10/076,7	
1 (48	, 1 2 7	004 gg			Applicant	SK Sharm	ıa, KB Ra	ınk		
B THE	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \				Filing Date 02/15/2002			Group	1641	
~	TRAS			U.S. PA	TENT DOCUM	ENTS				
Examiner Initial	-	Document Number Date			Name Class			Subclass Filing Date If Appropria		
	AA									
•	AB	ļ								
	AC									
	AD	 								
	AE	 		·					٠.	
	AF	 								
	AG	 								
	AH	ļ			<u> </u>					
	AI	<u></u>			· ·					
		Document	FOI Date	REIGN I	PATENT DOCU	MENTS				
		Number	Date		Country		Class	Subclass Translation		
	AJ								Yes	No
	AK									
	AL									
	AM									
	AN									+
			OTHER PRIOR A	RT (Inclu	ding Author, Title, D	ate, Pertinent Pa	ges, Etc.)			
	AO	OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.) Bancher, C. et al., "Accumulation of abnormally phosphorylated \u03c4 precedes the formation of neurofibrillary tangles in Alzheimer's disease," Brain Research 477(1-2):90-99 [1989];								
	AP	Bondareff, W. et al., "Molecular analysis of neurofibrillary degeneration in Alzheimer's Disease," American J. of Pathology 137(3):711-23 [1990];								
	AQ	Caceres, A. et al., "Inhibition of neurite polarity by tau antisense oligonucleotides in primary cerebellar neurons," Nature 343(6257):461-63 [1990];								
	AR	Drubin, D.G. et al., "Tau protein function in living cells," J. Cell Biology 103(6 PT 2):2739-46 [1986];								
	AS	Evans, D.B. et al., "Tau phosphorylation at Serine 396 and Serine 404 by human recombinant Tau protein kinase II inhibits Tau's ability to promote microtubule assembly," J. Biological Chemistry 275(32):24977-983 [2000];								
	AT	Glenner, G.G. et cerebrovascular	al., "Alzheimer's E amyloid protein," F)isease: l Biochemi	Initial report of t	he purificatio	n and cha	uracterizatio ications 120	n of a no)(3):885-9	vel 90

Aty. Docket No. 6322.N Applicant Applicant Applicant Applicant SK Sharma, KB Rank Filing Date 62/15/2002 Group Applicant AV Goodert, M. et al., "Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer diseases. Identification as the microtubule-associated protein tau." PNAS 85-4051-56 [1988]; AW Goodert, M. et al., "Multiple isoforms of human microtubule-associated protein tau." PNAS 85-4051-56 [1988]; AX Goodert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in neurofibrillary tangles of Alzheimer's Disease, Neuron 3(45:51-26 [1988]); AX Goodert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tabulin polymerization," The EMBIO Journal 8(13): 4225-30 [1980]; AZ Groenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct \(\tau\) proteins by polyacrylamide gel electrophoresis," PNAS 87(15:5827-31 [1990); BO Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family," Molecular and Cellular Biology 9(4):1388-96 [1980]; BP Horie, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6970):605-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TNS 14(6):218-19 [1991]; BC Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TNS 14(6):218-19 [1991]; BC Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TNS 14(6):218-19 [1991]; BC Kosik, K.S. et al., "Microtubule-associated protein tau (c) a major antigenic component of paired helical filaments in Alzheimer's microtubule-binding domains of tau: A model for self-association," Neuron 6(3):717-28 [1991]; BC Lee, V.MY., et al., "Tae files	·	<u> </u>							
Applicant SK Sharma, KB Rank Filing Date 02/15/2002 Group Applicant SK Sharma, KB Rank Filing Date 02/15/2002 Group Applicant SK Sharma, KB Rank Filing Date 02/15/2002 Group Average of Alzheimer disease: Identification as the microtubule-associated protein tau, "PNAS 85-4051-55 [1988]; AW Goedert, M. et al., "Multiple isoforms of human microtubule-associated protein tau," PNAS 85-4051-55 [1988]; AX Goedert, M. et al., "Multiple isoforms of human microtubule-associated protein tau," PNAS 85-4051-55 [1988]; AX Goedert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization," The EMBO Journal 9(13), 4225-30 [1980]; AY Goedert, Michael, "Tau protein and the neurofibrillary pathology of Alzheimer's Disease," Trends in Neuroscience 18(11):460-65 [1993]; AZ Greenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct \(\tau\) proteins by polyacrylamide gel electrophoresis," PNAS 97(15):5827-31 [1990]; BO Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family, Molecular and Cellular Biology 94(1398-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microbuble-associated protein tau (c) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(1):404-44 [1986]; BY Kosik, K.S. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):255-88 [1988]; BV Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Turnet of inormal tau," Science 221(494):675-78 [1991]; BW Lee, V.MY., et al., "The				Atty	Docket No.	Serial No.			
Applicant SK Sharma, KB Rank Filing Date 02/15/2002 Group Applicant SK Sharma, KB Rank Filing Date 02/15/2002 Group AV Goedert, M. et al., "Cloning and sequencing of the cDNA encoding a core protein of the paired belical filament of Alzheimer disease: Identification as the microtubule-associated protein tau." PNAS 85:4051-55 [1988]; AW Goedert, M. et al., "Multiple isoforms of human microtubule-associated protein tau: Sequences and localization in neurofibrillary tangles of Alzheimer's Disease," Neuron 3(4):519-26 [1989]; AX Goedert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization," The EMBO Journal 9(13): 4225-30 [1980]; AY Goedert, Michael, "Tau protein and the neurofibrillary pathology of Alzheimer's Disease," Trends in Neuroscience 15(11):460-65 [1993]; AZ Greenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct \(\tau\) proteins by polyacrylamide gel electrophoresis," PNAS 97(15):5827-31 [1990]; BO Himmler, Adolf, 'Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family, Molecular and Cellular Biology 9(4):1388-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):685-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BS Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S., et al., "Microtubule-associated protein tau (c) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):404-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau." A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, V.M.Y., et al., "A88: A major subunit of paired helical filaments and derivatized forms of nor					6322.N	10/076,708			
Filing Date 02/15/2002 Group AV Goedert, M. et al., "Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer disease: Identification as the microtubule-associated protein tau," PNAS 85-4051-55 [1986]; AW Goedert, M. et al., "Multiple isoforms of human microtubule-associated protein tau: Sequences and localization in neurofibrillary tangles of Alzheimer's Disease," Neurosciated protein tau: Sequences and localization in neurofibrillary tangles of Alzheimer's Disease, Neurosciated protein: correlation with the tau pattern in brain and effects on tubulin polymerization," The EMBO Journal 9(13): 4225-30 [1990]; AY Goedert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization," The EMBO Journal 9(13): 4225-30 [1990]; AZ Greenberg, S.C., et al., "A preparation of Alzheimer paired helical filaments that displays distinct \(\tau\) proteins by polyacrylamide gel electrophoresis, "PNAS 87(15):5827-31 [1990]; BO Himmler, Adolf, "Structure of the bovine tau spen: Alternatively spliced transcripts generate a protein family," Molecular and Cellular Biology 9(4):1389-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 21(6070):660-7 (1986); BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 (1990); RR Kosik, K.S., *Alzheimer plaques and tangles: advances on both fronts," TTNS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (r) is a major antigenic component of paired helical filaments in Alzheimer diseases," PNAS 83(11):4044-48 [1986]; BT Keiezk Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau	LIST OF REFERENCES CITED BY APPLICANT (Usersweral sheets if necessary)								
Group Group Group Group Group Group Group Group Godert, M. et al., "Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer disease. Identification as the microtubule-associated protein tau." PNAS 85:4051-55 [1988]; Godert, M. et al., "Multiple isoforms of human microtubule-associated protein tau: Sequences and localization in neurofibrillary tangles of Alzheimer's Disease," Neuron 3(4):519-26 [1989]; AX Goedert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization," The EMBO Journal 9(31): 4225-30 [1990]; AY Goedert, Michael, "Tau protein and the neurofibrillary pathology of Alzheimer's Disease," Trends in Neuroscience 16(11):460-65 [1989]; AZ Groenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct \(\tau\) proteins by polyacrylamide gel electrophoresis," PNAS 67(15):5627-31 [1990]; BO Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family," Molecular and Cellular Biology 9(4):1389-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's diseases," Current Opinion in Cell Biology 2(1):101-4 [1990]; Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; Kosik, K.S., et al., "Microtubule-associated protein tau (v) is a major antigenic component of paired helical filaments in Alzheimer diseases," PNAS 83(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(3):717-28 [1991]; BU Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BY Spillantini, M.G. et al., "Tau p	2 2004			Applicant	SK Sharma, KB Rar	ık			
Goedert, M. et al., "Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer disease: Identification as the microtubule-associated protein tau," PNAS 85:4051-55 [1988]; AW Goedert, M. et al., "Multiple isoforms of human microtubule-associated protein tau: Sequences and localization in neurofibrillary tangles of Alzheimer's Disease," Neuron 3(4):519-26 [1989]; AX Goedert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization," The EMBO Journal 9(13): 4225-30 [1990]; AY Goedert, Michael, "Tau protein and the neurofibrillary pathology of Alzheimer's Disease," Trends in Neuroscience 16(11):460-66 [1993]; AZ Greenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct \(\tau\) proteins by polyacrylamide gel electrophoresis," PNAS 87(15):5827-31 [1990]; BO Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family, Molecular and Cellular Biology 9(3):1389-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):805-7 [1986]; BR Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (c) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):3044-46 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(6):717-26 [1991]; BU Lee, C. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1986]; BY Lee, V.M. Y., et al., "A68: A major subunit of paired helical filaments and derivatized forms	1 \			Filing Date	02/15/2002	Group			
AV Goedert, M. et al., "Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer disease: Identification as the microtubule-associated protein tau." PNAS 85:4051-55 [1988]; AW Goedert, M. et al., "Buttiple isoforms of human microtubule-associated protein tau: Sequences and localization in neurofibrillary tangles of Alzheimer's Disease," Neuron 3(4):519-26 [1989]; AX Goedert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization," The EMBO Journal 9(13): 4225-30 [1990]; AY Goedert, Michael, "Tau protein and the neurofibrillary pathology of Alzheimer's Disease," Trends in Neuroscience 16(11):460-65 [1993]; AZ Greenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct \(\tau\) proteins by polyacrylamide gel electrophoresis," PNAS 87(15):5827-31 [1990]; BO Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family, Molecular and Cellular Biology 9(3):1389-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986]; BR Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (c) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):404-44 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-26 [1991]; BU Lee, V.MY., et al., "Afs: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 23(4834):285-88 [1988]; BY Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's	- 3	TRAN	[1984];						
AW Goedert, M. et al., "Multiple iseforms of human microtubule-associated protein tau: Sequences and localization in neurofibrillary tangles of Alzheimer's Disease," Neuron 3(3):519-26 [1989]; AX Goedert, M. et al., "Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization," The EMBO Journal 9(13): 4225-30 [1990]; AY Goedert, Michael, "Tau protein and the neurofibrillary pathology of Alzheimer's Disease," Trends in Neuroscience 16(11):460-65 [1983]; AZ Greenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct τ proteins by polyacrylamide gel electrophoresis," PNAS 87(15):5627-31 [1990]; BO Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family," Molecular and Cellular Biology 9(3):1389-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; Kosik, K.S. et al., "Microtubule-associated protein tau (τ) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, Q. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):633-56 [1992]; BY Spillantini, M.G. et al., "Tau p			Goedert, M. et al., "Cloning and sequencing of the cDNA encoding a company to the project of the						
AY Goedert, Michael, "Tau protein and the neurofibrillary pathology of Alzheimer's Disease," Trends in Neuroscience 16(11):460-65 [1993]; AZ Greenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct τ proteins by polyacrylamide gel electrophoresis," PNAS 87(15):5827-31 [1990]; Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family," Molecular and Cellular Biology 9(4):1389-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (t) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 33(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of faux a model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(6):1858-62 [1975].		AW	Goedert, M. et al., "Multiple isoforms of	human microtub	ule-associated protein to				
AZ Greenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct τ proteins by polyacrylamide gel electrophoresis." PNAS 87(15):5827-31 (1990); BO Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family," Molecular and Cellular Biology 9(4):1389-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (τ) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):4044-48 [1986]. BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):225-38 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		AX	Goedert, M. et al., "Expression of separate isoforms of human tau protein; correlation with the description of the separate isoforms of human tau protein; correlation with the description of the separate isoforms of human tau protein; correlation with the description of the separate isoforms of human tau protein; correlation with the separate isoforms of human tau protein wit is separate isoforms of human tau protein with the separate iso						
BO Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family," Molecular and Cellular Biology 9(4):1389-96 [1989]; BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (r) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "468: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		AY	Goedert, Michael, "Tau protein and the Neuroscience 16(11):460-65 [1993];	neurofibrillary pa	athology of Alzheimer's I	Disease," Trends in			
BP Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986]; BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (r) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-66 [1992]; Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		AZ	Greenberg, S.G., et al., "A preparation of Alzheimer paired helical filaments that displays distinct τ proteins by polyacrylamide gel electrophoresis," PNAS 87(15):5827-31 [1990]:						
BQ Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 [1990]; BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (t) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		ВО	Himmler, Adolf, "Structure of the bovine tau gene: Alternatively spliced transcripts generate a protein family."						
BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (\tau) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		BP	Horio, T. et al., "Visualization of the dynamic instability of individual microtubules by dark-field microscopy," Nature 321(6070):605-7 [1986];						
BR Kosik, K.S., "Alzheimer plaques and tangles: advances on both fronts," TINS 14(6):218-19 [1991]; BS Kosik, K.S. et al., "Microtubule-associated protein tau (\tau) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		BQ	Kosik, K.S., "Tau protein and Alzheimer's disease," Current Opinion in Cell Biology 2(1):101-4 (1990).						
BS Kosik, K.S. et al., "Microtubule-associated protein tau (t) is a major antigenic component of paired helical filaments in Alzheimer disease," PNAS 83(11):4044-48 [1986]; BT Ksiezak-Reding et al., "Structural stability of paired helical filaments requires microtubule-binding domains of tau: A model for self-association," Neuron 6(5):717-28 [1991]; BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		BR							
BU Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse brain," Science 239(4837):285-88 [1988]; BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		BS	Kosik, K.S. et al., "Microtubule-associated protein tau (7) is a major antigenic component of pointed believed						
BV Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991]; BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975].		BT	Ksiezak-Reding et al., "Structural stability of paired helical filaments requires migrotubule hinding demoires						
BW Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992]; BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975]. CO CP		BU	Lee, G. et al., "The primary structure and heterogeneity of tau protein from mouse havin." Science						
BX Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998]; BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975]. CO CP		BV	Lee, V.MY., et al., "A68: A major subunit of paired helical filaments and derivatized forms of normal tau," Science 251(4994):675-78 [1991];						
BY Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975]. CO CP		BW	Lee, V.MY., et al., "The disordered neuronal cytoskeleton in Alzheimer's disease," Current Opinion in Neurobiology 2(5):653-56 [1992];						
21(10):428-33 [1998]; BZ Weingarten, M.D. et al., "A protein factor essential for microtubule assembly," PNAS 72(5):1858-62 [1975]. CO CP		ВХ	Mailliot, C. et al., "Phosphorylation of specific sets of tau isoforms reflects different neurofibrillary degeneration processes," FEBS Letters 433(3):201-04 [1998];						
CO CP		BY	Spillantini, M.G. et al., "Tau protein pathology in neurodegenerative diseases," Trends in Neuroscience 21(10):428-33 [1998];						
CO CP		BZ	Weingarten, M.D. et al., "A protein factor essential for microtubule assembly " PNAS 72(5):1859 69 (1075)						
		СО	17.00-02 [1970].						
Examiner Date Considered		СР							
	Examiner			Date Considered					